

Errata: Evaluation of garlic extracts antifungal potential against causative fungi of dermatomycosis in Chile

Errata: Evaluación del potencial antifúngico de extractos de ajos frente a hongos causantes de dermatomicosis en Chile

Peggy Vieille Oyarzo^{1,*} , Melissa Noguera Gahona¹ , Rodrigo Cruz Choappa² 

Correction to the article: Vieille Oyarzo, P., Noguera Gahona, M., Cruz Choappa, R., & Cogliati, M. (2024). Evaluation of garlic extracts antifungal potential against causative fungi of dermatomycosis in Chile. *ARS Medica*, 49(3), 38–43. <https://doi.org/10.11565/arsmed.v49i3.2075>

This correction applies to the Materials and Methods, Results, and Discussion sections, as well as to Table 1, Figure 1, and Figure 2. For completeness and transparency, the previous versions of Table 1, Figure 1, and Figure 2 (incorrect versions) are presented below, as well as the corrected Table 1, Figure 1, and Figure 2.

In the original article, the unit reported as mg/L, should have been reported is g/L.

The original article has also been corrected.

Incorrect Table 1: Antifungal activity of the three garlic extracts tested in the present study.

Extract concentration (mg/L)	Chinese fresh garlic		Chilote fresh garlic		Chilote black garlic	
	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>
0.8	+++	+++	+++	-	+++	+++
1.6	+++	-	-	-	+++	+
3.2	+++	-	-	-	+++	-
6.4	+++	-	-	-	-	-

+++ , ++ , + : growth compared to the positive control; - : absence of growth

Corrected Table 1: Antifungal activity of the three garlic extracts tested in the present study.

Extract concentration (g/L)	Chinese fresh garlic		Chilote fresh garlic		Chilote black garlic	
	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>
0.8	+++	+++	+++	-	+++	+++
1.6	+++	-	-	-	+++	+
3.2	+++	-	-	-	+++	-
6.4	+++	-	-	-	-	-


+++ , ++ , + : growth compared to the positive control; - : absence of growth

(1) Facultad de Medicina, Universidad de Valparaíso. Viña del Mar, Chile.


(2) Centro de Diagnóstico e Investigación de Enfermedades Infecciosas (CDIEI), Universidad de Valparaíso. Viña del Mar, Chile.

*Corresponding author: peggy.vieille@uv.cl







Incorrect Figure 1: Growth of *Candida albicans* in the three types of garlic compared to the positive control.



Incorrect Figure 2: Growth of *Trichophyton rubrum* in the three types of garlic compared to the positive control.



Corrected Figure 1: Growth of *Candida albicans* in the three types of garlic compared to the positive control.



Corrected Figure 2: Growth of *Trichophyton rubrum* in the three types of garlic compared to the positive control.